

# Hooded skunk

## Morphology

It can be distinguished from the similar [striped skunk](#) (*M. mephitis*) by its longer tail and longer, much softer coat of fur, and larger tympanic bullae.<sup>[2]</sup> A ruff of white fur around its neck gives the animal its common name. Three color phases are known and in all three, a thin white medial stripe is present between the eyes: black-backed with two lateral white stripes, white-backed with one dorsal white stripe, or entirely black with a few white hairs in the tail.<sup>[3][4]</sup>

## Ecology

The hooded skunk ranges from the [Southwestern United States](#) to [southern Mexico](#), but is most abundant in [Mexico](#). These skunks are found to be 50% or less smaller in size in southern Mexico than in the Southwestern United States.<sup>[5]</sup> It is found in [grasslands](#), [deserts](#), and in the foothills of [mountains](#), avoiding high elevations. It tends to live near a water source, such as a river. The females tend to be 15% smaller in size than the males<sup>[6]</sup> and their breeding season is between February and March.<sup>[4]</sup> The litter size ranges from three to eight.<sup>[7]</sup>

## Diet

The diet of the hooded skunk consists mostly of vegetation, especially prickly pear (*Opuntia* spp.), but it will readily consume [insects](#), small [vertebrates](#), and bird eggs <sup>[4]</sup> as well. No cases of rabies are reported,<sup>[8][9]</sup> but they host a range of parasites, including nematodes, roundworms, and fleas.<sup>[4]</sup>

## Behavior

Hooded skunks are solitary, but they might interact at a feeding ground without showing any signs of aggression.<sup>[10]</sup> They shelter in a burrow or a nest of thick plant cover during the day and are active at night. Like *M. mephitis*, for self-defense, they spray volatile components from their anal glands.<sup>[11]</sup>

## Characteristics

Hooded skunks are currently not endangered. They are very abundant in Mexico and can live in human suburban areas mostly on pastures and cultivated fields.<sup>[12]</sup> Their fur has low economic value.<sup>[7]</sup> However, their fat<sup>[11]</sup> and scent glands<sup>[10]</sup> can be used for medicinal purposes. In some parts of their range, their flesh is considered a delicacy.<sup>[13]</sup> Other common names for the hooded skunk include: *mofeta rayada* (Spanish), *moufette à capuchon* (French), *pay* (Maya), southern skunk, white-sided skunk, and *zorillo*.<sup>[14]</sup>

## External links

- [Smithsonian Institution - North American Mammals: \*Mephitis macroura\*](#)

## References

- ↑  *Template:IUCN2008*
- ↑ Hall, E. R. (1981). *The mammals of North America*. Second edition. New York: John Wiley & Sons. pp. 601–1181.
- ↑ Hoffmeister, D. F. (1986). *Mammals of Arizona*. Tucson: University of Arizona Press.
- ↑ <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> Patton, R. F. (1974). *Ecological and behavioral relationships of the skunks of Trans Pecos Texas*. Ph.D. dissertation. Texas A&M University. p. 199.
- ↑ Janzen, D. H. and W. Hallwachs (1982). *The hooded skunk, Mephitis macroura, in lowland northwestern Costa Rica*. *Brenesia*. pp. 19/20:549–552.
- ↑ Rosatte, R. C. (1987). *Striped, spotted, hooded, and hog-nosed skunk*. Toronto, Canada: Ontario Ministry of Natural Resources.
- ↑ <sup>a</sup> <sup>b</sup> Bailey, V. (1932). "Mammals of New Mexico". *North American Fauna* (53): 1–412.
- ↑ Aranda, M.; L. Lopez-De Buen (1999). "Rabies in skunks from Mexico". *Journal of Wildlife Diseases* (35): 574–577.
- ↑ Ceballos, G., And A. Miranda (1986). *Los mamíferos de Chmela, Jalisco: manual de campo*. Mexico City, Mexico: Universidad Nacional Autonoma de Mexico.
- ↑ <sup>a</sup> <sup>b</sup> Reid, F. A. (1997). *A field guide to the mammals of Central America and south east Mexico*. New York: Oxford University Press.
- ↑ <sup>a</sup> <sup>b</sup> Dalquest, W. W. (1953). *Mammals of the Mexican state of San Luis Potosi*. Balton Rouge: Louisiana State University Press.
- ↑ Yeen, Ten Hwang; Serge Lariviere (26 December 2001). "Mephitis macroura". *Mammalian Species* (686): 1–3.
- ↑ Davis, W. B. (1944). "Notes on Mexican mammals". *Journal of Mammalogy* (25): 370–402.
- ↑ Borror, D. J. (1960). *Dictionary of word roots and combining forms*. Palo Alto, California: National Press Books.

- ↑ Irwin, M. 2001. *Mephitis macroura* (on-line), [Animal Diversity Web](#). Accessed September 13, 2006